



PLACEMENT®

DRIFT CONTROL AGENT & DEPOSITION AID

PRINCIPAL FUNCTIONING AGENTS:

Petroleum distillates, amine salts of fatty acids, aromatic acid 100%

Contains petroleum distillates. All ingredients are exempt for the requirements of a tolerance in 40 CFR 180.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear chemical-resistant gloves when mixing or handling concentrate. Wash thoroughly with soap and water after handling.

First Aid -- If Swallowed: Immediately call a Poison Control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. If on Skin or Clothing: Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE: Store in original container only and keep sealed. Store in closed storage areas.

ADJUVANT DISPOSAL: Wastes resulting from the use of this product may be disposed of through on-site spray application or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent) during mixing and loading. Recycling decontaminated containers is the best option of container disposal. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. Contact your state and local ACRC recycler or visit the ACRC web page at www.acrecycle.org. Decontaminated containers may also be disposed of in a sanitary landfill.

GENERAL INFORMATION

PLACEMENT is a spray adjuvant designed to maximize pesticide performance by improving deposition of the spray application on to the intended target. PLACEMENT improves coverage and reduces evaporation of pesticides being applied by ground or air. PLACEMENT encapsulates pesticides into positively charged uniform capsules. These capsules are attracted to negatively charged plant tissue and/or soil particles resulting in maximized on-target delivery of spray solution.

PLACEMENT is unique and does not affect or change spray droplet size like other drift control adjuvants. Therefore, spray droplet size can be managed by proper selection of nozzle type, size and pressure. Along with good judgment and the proper selection and use of spray equipment, PLACEMENT can help professional applicators ensure the maximum performance of pesticides.

WA Reg. No. 1381-02003

Distributed by:
Agrilience, LLC
P.O. Box 64089, St. Paul, MN 55164-0089

NET CONTENTS
0/H29/5

DIRECTIONS FOR USE FOR TERRESTRIAL USE ONLY – NOT FOR AQUATIC USE

Prior to use, read all directions on this label.

The proper rate of PLACEMENT[®] is dependent upon the type of pesticide formulation being applied. First, determine the pesticide formulation being applied, such as liquid, flowable, or powder. Then select the proper rate according to the following table:

RATE OF PLACEMENT [®]	Pesticide formulation* and use rate
4 ounces	Per 1 pint (16 oz) of E.C. or A. S. – liquid formulations
2 ounces	Per 1 pound of D.F. or W. P. – dry formulations
2 ounces	Per 1 quart (32 oz) of L.F. – liquid flowables

*E.C. = Emulsifiable Concentrates such as Trust[®], A.S.= Aqueous Solution such as Roundup[®], D.F.= Dry Flowables such as Accent[®], W.P.= Wettable Powders such as Connect[™] 20 WSP, L.F.= Liquid Flowables such as Sterling[®] Plus.

Note: The above use rates equate to 1 ounce of PLACEMENT[®] for every 4 ounces of E.C. or A.S. herbicide. For example, when using Roundup[®] Weather Max[™], Roundup[®] Original[™], Roundup Original Max, Roundup Ultra Max[®] II, Touchdown[®] IQ, or Cornerstone[®], use 4 oz of PLACEMENT[®] for 16 oz of the herbicide and 8 oz. of PLACEMENT[®] for 32 oz of herbicide.

DO NOT ADD THIS PRODUCT AT A RATE THAT EXCEEDS 5% OF THE FINISHED SPRAY VOLUME.

Use with Pyrethroid Insecticides – Use PLACEMENT[®] at a 1:1 ratio. Example: 3 oz of pyrethroid per acre + 3 oz of PLACEMENT[®] per acre.

MIXING: Add PLACEMENT[®] after pesticides or other adjuvants are added but prior to complete filling of the spray tank. For maximum drift control when using Emulsifiable Concentrates or Aqueous Solutions, pre-mixing of PLACEMENT[®] and the EC or AS or the use of a chemical inductor is suggested.

When mixing with Flowables (dry or liquid) and Wettable Powders, add flowables or wettable powders to the half filled spray tank and agitate thoroughly while continuing to fill spray tank. Then add the PLACEMENT[®] last while continuing agitation.

Notice of Warranty

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.

AgriSolutions[™] is a trademark and Cornerstone[®], Placement[®], Sterling[®], and Trust[®] are registered trademarks of Agrilience, LLC.

Connect[™] is a trademark of Bayer.

Accent[®] is a registered trademark of E.I. duPont de Nemours and Company.

Original[™] and Weather Max[™] are trademarks and Roundup[®] and UltraMax[®] are registered trademarks of Monsanto Technology LLC.

Touchdown[®] is a registered trademark of Syngenta.